

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/82904 A1

- (51) International Patent Classification⁷: A61K 9/32. (74) Agent: ROSENSTOCK, Jerome; Frommer Lawrence & Haug LLP, 745 Fifth Avenue, New York, NY 10151 (US).
9/34, 9/36, 9/38
- (21) International Application Number: PCT/US01/00039 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 2 January 2001 (02.01.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/201,233 1 May 2000 (01.05.2000) US
09/703,023 31 October 2000 (31.10.2000) US
- (71) Applicant: AEROPHARM TECHNOLOGY, INC. [US/US]; Raritan Center, Campus Plaza, 18 Mayfield Avenue, Edison, NJ 08837 (US).
- (72) Inventors: CUTIE, Anthony, J.; 10 Tillman Court, Bridgewater, NJ 08807 (US). ADJEI, Akwete, L.; 15 Tillman Court, Bridgewater, NJ 08807 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/82904 A1

(54) Title: A CORE FORMULATION

(57) Abstract: This invention relates to a combination drug product comprising troglitazone, e.g. its hydrochloride, and a biguanide, e.g. metformin. In particular, the product comprises a core of metformin, at least a portion thereof has a layer or coat thereon of troglitazone.

A CORE FORMULATION

This application claims priority from U.S. provisional application Serial No. 60/201,233 filed May 1, 2000, which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

Field of the Invention

This invention relates to a core formulation, and, more particularly, to a core formulation comprising a first layer comprising troglitazone, which covers at least a portion of a core comprising a biguanide, such as for example metformin (i.e., glucophage).

Description of the Related Art

Metformin and troglitazone, and their salts such as the hydrochlorides, maleates, tartrates, etc., are two active ingredients of anti-diabetic drugs that are used to treat diabetic patients, e.g. human beings. These two active agents are administered orally to patients in need thereof in protocols calling for the single administration of either ingredient. Heretofore, there has not been revealed or hinted at combining both ingredients and certainly not a physically combined core formulation comprising both ingredients. The advantage of such a core formulation is advantageous to patients and prescribers because both medicaments are synergistic to each other in the body when used in the management of blood glucose control, i.e., diabetes.

SUMMARY OF THE INVENTION

This invention relates to a core formulation, and, more particularly, to a core formulation comprising a first layer comprising troglitazone or a derivative thereof, e.g. troglitazone hydrochloride, which covers at least a portion of a core comprising a biguanide. A typical biguanide is metformin. It typically is used clinically as a pharmaceutically acceptable salt, preferably the hydrochloride salt. A commercial form of metformin hydrochloride is available as glucophage. Its chemical name is N,N-dimethylimidodicarbonimidic diamide hydrochloride. Metformin hydrochloride is a hydrochloride salt of metformin base, and as used herein, "metformin" means the base compound as well as its pharmaceutically acceptable salts. Metformin is used clinically to manage non-insulin dependent diabetes mellitus ("NIDDM"), particularly in patients who are not effectively treated

with a sulfonylurea. While it is not chemically related to the sulfonylureas, it is routinely utilized in combination with a sulfonylurea, and has been shown to be synergistic in some cases. Other biguanides such as phenformin, buformin etc. can also be used. Additionally, in the treatment of a diabetic patient the metformin and
5 the troglitazone, e.g. its hydrochloride, are present in effective amounts to provide such treatment.

DETAILED DESCRIPTION OF THE INVENTION

Metformin is an active ingredient for a commercially available drug employed to treat diabetes mellitus in a host or mammal, e.g. a human being,
10 another animal. The typical daily effective dose for the oral treatment of a mammal, i.e., a human, ranges from 500 mg to 2550 mg. Typically, the dose is a single dose of 500 mg to 850 mg.

Troglitazone hydrochloride, (Rezulin®), is an active ingredient for a commercially available drug employed to treat diabetes mellitus in a host, e.g. a
15 human being. The typical daily effective dose for the oral administration to a mammal, e.g. a human being, ranges from 200 mg to 400 mg, given as a single dose.

Heretofore, the metformin and pioglitazone hydrochloride have not been administered together to try to improve the effectiveness of either drug, although co-administration of the two has been proposed [*Whitcomb; et al., United States Patent No. 6,011,049*]. However, a combined form of the drugs, i.e. a single
20 integral unit thereof has not heretofore been reported. The present invention provides such a single integral unit in the form of a core formulation.

As indicated above, the relative concentrations of each drug is such that a first layer comprising metformin is prepared. The first layer covers at least a
25 portion of a core comprising metformin. Depending upon the rate of administration and metabolism of the patient destined to be treated, and the concentrations of each ingredient desired for each drug, the first layer may cover only a portion of the core or encompass the entire core. For example, one quarter of the core to about three fourths of the tablet core. The first layer should comprise troglitazone hydrochloride
30 because its dose requirement is lower compared to metformin. Additionally, it is slightly non-polar, its solubility rate slower, and its absorption rate thus dependent

on its dissolution rate in the contents of the gastrointestinal tract compared with metformin.

It is to be understood, depending upon the rate of administration to the patient desired, either the first layer or the core may contain a mixture of the two active ingredients or both the first layer and the core may contain the two active ingredients with different and varying concentrations of one or both active ingredients.

The first layer of the core comprises troglitazone, e.g. its hydrochloride, in an amount of 0.01 % to 20% of the total weight of the core formulation, whereas, the metformin in the core is present in an amount of 10% to 97.5% of the total weight of the core formulation.

Where combinations of the two active ingredients are present in the first layer and/or the core, the amounts of troglitazone, e.g. its hydrochloride, ranges from 1 mg to 45 mg whereas the metformin ranges from 100 mg to 2550 mg.

Finally, it is to be understood that a third pharmacologically active material, e.g. a drug, such as for example sulfonylureas, α -glucosidase inhibitors, meglitinides, and ACE inhibitors can admixed with the active ingredients in the first layer and/or the core.

The α -glucosidase inhibitors [Jean-Bernard Ducep et al., US Patent No. 5,504,078], bisglucosylmoranoline derivatives [UK Patent No. GB 2 088 365 A], and glucosylmoranoline derivatives [European Patent No. 87112480.6] include the following medicaments: 1.5-Dideoxy-4-O(α ,D-glucopyranosyl)-1,5-[6,7-dideoxy-7-D-glucoheptopyranosyl]imino]-D-glucitol; 1.5-Dideoxy-4-O(α ,D-glucopyranosyl)-1,5-[(1-deoxy-D-fructofuranosyl)imino]-D-glucitol; 1.5-Dideoxy-4-O(α ,D-glucopyranosyl)-1,5-[(4-deoxy-4-D-glucopyranosyl)imino]-D-glucitol; 1.5-Dideoxy-4-O(α ,D-glucopyranosyl)-N-[6-deoxy-1-(6-O-D-glucopyranosyl)- α -D-glucopyranosyl]-1, 5-imino-D-glucitol; 1.5-Dideoxy-4-O(α ,D-glucopyranosyl)-N-[6,7-dideoxy-1-(6-O-D-glucopyranosyl)-7- α -D -glucoheptopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-4-O(α ,D-glucopyranosyl)-1,5-[(4-deoxy-4-D-glucopyranosyl)methylimino]-D-glucitol; 1.5-Dideoxy-4-O(α ,D-glucopyranosyl)-N-[4-deoxy-1-(4-O-D-glucopyranosyl)- α -D-glucopyranosyl]-

1,5-imino-D-glucitol; 1.5-Dideoxy-4-O(.alpha.,D-glucopyranosyl)-1,5-{{2(1-D-
 arabinofuranose)ethyl}imino}-D-glucitol; 1.5-Dideoxy-4-O(.alpha.,D-
 glucopyranosyl)-N-[4-deoxy-1-(6-O-D-glucopyranosyl)-.alpha.-D-glucopyranosyl]-
 1,5-imino-D-glucitol; 1.5-Dideoxy-4-O(.alpha.,D-glucopyranosyl)-N-{{4-deoxy-1-
 5 (4-O-D-glucopyranosyl)-4-.alpha.-D -glucopyranosyl)methyl}-1,5-imino-D-glucitol;
 1.5-Dideoxy-4-O(.alpha.,D-glucopyranosyl)-N-{{4-deoxy-1-(6-O-D-
 glucopyranosyl)-4-.alpha.-D -glucopyranosyl)methyl}-1,5-imino-D-glucitol; 1.5-
 Dideoxy-4-O(.alpha.,D-glucopyranosyl)-1,5-[(6-deoxy-1-O-methyl-6-β-D-
 glucopyranosyl)-imino-D -glucitol; 1.5-Dideoxy-4-O(.alpha.,D-glucopyranosyl)-
 10 1,5-[(6,7-dideoxy-1-O-methyl-7-β-D-glucoheptopyranosyl) imino]-D-glucitol; 1.5-
 Dideoxy-4-O(.alpha.,D-glucopyranosyl)-1,5-[(1-deoxy-2-O-methyl-β-D-
 fructofuranosyl)imino]-D-glucitol, 1.5-Dideoxy-4-O(.alpha.,D-glucopyranosyl)-1,5-
 [(4-deoxy-1-O-methyl-4-β-D-glucopyranosyl)imino]-D-glucitol; 1.5-Dideoxy-4-
 O(.alpha.,D-glucopyranosyl)-N-[6-deoxy-1-(1-O-methyl-6-O-β-D-glucopyranosyl)-
 15 .alpha.-D -glucopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-4-O(.alpha.,D-
 glucopyranosyl)-N-[6,7-dideoxy-1-(1-O-methyl-6-O-β-D-glucopyranosyl)-7-.alpha.-
 D-glucoheptopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-4-O(.alpha.,D-
 glucopyranosyl)-1,5-[(4-deoxy-1-O-methyl-4-β-D-glucopyranosyl)methylimino]-D-
 glucitol; 1.5-Dideoxy-4-O(.alpha.,D-glucopyranosyl)-N-[4-deoxy-1-(1-O-methyl-4-
 20 O-8-D-glucopyranosyl)-.alpha.-D -glucopyranosyl]-1,5-imino-D-glucitol; 1.5-
 Dideoxy-4-O(.alpha.,D-glucopyranosyl)-1,5-{{2-(1-O-methyl-1-β-D-
 arabinofuranosyl)ethyl}imino}-D-glucitol; 1.5-Dideoxy-4-O(.alpha.,D-
 glucopyranosyl)-N-[4-deoxy-1-(1-O-methyl-6-O-β-D-glucopyranosyl)-.alpha.-D-
 glucopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-4-O(.alpha.,D-glucopyranosyl)-
 25 N-{{4-deoxy-1-(1-O-methyl-4-O-β-D-glucopyranosyl)-4-.alpha.-D-
 glucopyranosyl)methyl}-1,5-imino-D-glucitol; 1.5-Dideoxy-4-O(.alpha.,D-
 glucopyranosyl)-N-{{4-deoxy-1-(1-O-methyl-6-O-β-D-glucopyranosyl)-4-.alpha.-D-
 glucopyranosyl)methyl}-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-
 glucopyranosyl)-1,5-[6,7-dideoxy-7-D-glucoheptopyranosyl)imino]-D-glucitol; 1.5-
 30 Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-[(1-deoxy-D-fructofuranosyl)imino]-D-
 glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-[(4-deoxy-4-D-
 glucopyranosyl)imino]-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-

- [6-deoxy-1-(6-O-D-glucopyranosyl)-.alpha.-D-glucopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-[6,7-dideoxy-1-(6-O-D-glucopyranosyl)-7-.alpha.-D -glucoheptopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-[(4-deoxy-4-D-glucopyranosyl)methylimino]-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-[4-deoxy-1-(4-O-D-glucopyranosyl)-.alpha.-D-glucopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-{[2(1-D-arabinofuranose)ethyl]imino}-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-[4-deoxy-1-(6-O-D-glucopyranosyl)-.alpha.-D-glucopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-{[4-deoxy-1-(4-O-D-glucopyranosyl)-4-.alpha.-D -glucopyranosyl]methyl}-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-{[4-deoxy-1-(6-O-D-glucopyranosyl)-4-.alpha.-D -glucopyranosyl]methyl}-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-[(6-deoxy-1-O-methyl-6-β-D-glucopyranosyl) -imino]-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-[(6,7-dideoxy-1-O-methyl-7-β-D-glucoheptopyranosyl) imino]-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-[(1-deoxy-2-O-methyl-β-D-fructofuranosyl)imino]-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-[(4-deoxy-1-O-methyl-4-β-D-glucopyranosyl)imino]-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-[6-deoxy-1-(1-O-methyl-6-O-β-D-glucopyranosyl)-.alpha.-D -glucopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-[6,7-dideoxy-1-(1-O-methyl-6-O-β-D-glucopyranosyl) -7-.alpha.-D-glucoheptopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-[(4-deoxy-1-O-methyl-4-β-D-glucopyranosyl)methylimino]-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-[4-deoxy-1-(1-O-methyl-4-O-β-D-glucopyranosyl)-.alpha.-D -glucopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-1,5-{[2-(1-O-methyl-1-β-D-arabinofuranosyl)ethyl]imino}-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-[4-deoxy-1-(1-O-methyl-6-O-β-D-glucopyranosyl)-.alpha.-D -glucopyranosyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-glucopyranosyl)-N-[(4-deoxy-1-(1-O-methyl-4-O-β-D-glucopyranosyl)-4-.alpha.-D-glucopyranosyl)methyl]-1,5-imino-D-glucitol; 1.5-Dideoxy-6-O(.alpha.,D-

glucopyranosyl)-N-([4-deoxy-1-(1-O-methyl-6-O- β -D-glucopyranosyl)-4- α -D-glucopyranosyl]methyl)-1,5-imino-D-glucitol.

The list of medicaments includes acid addition salt forms with such inorganic acids such as, for example, hydrochloric, hydrobromic, sulfuric, phosphoric and like acids; with organic carboxylic acids such as, for example, acetic, propionic, glycolic, lactic, pyruvic, malonic, succinic, fumaric, maleic, tartaric, citric, ascorbic, maleic, hydroxymaleic and dihydroxymaleic, benzoic, 2-acetoxybenzoic, mandelic and like acids; and with organic sulfonic acids such as methanesulfonic acid and p-toluenesulfonic acid.

The sulfonylureas are a class of compounds that have been widely employed to treat diabetes. Such compounds are well known, for example, as described in U.S. Pat. Nos. 3,454,635; 3,669,966; 2,968,158; 3,501,495; 3,708,486; 3,668,215; 3,654,357; and 3,097,242. Especially preferred sulfonylureas to be employed in the combinations of this invention are glyburide, glipizide, tolbutamide, tolazamide, glisoxepid, chlorpropamide, glibornuride, glimepiride, phenbutamide, and tolcyclamide. Other medicaments, such as, for example, an antibiotic, a vitamin, a drug that works on the heart, or in the liver, may be admixed with the active ingredients in the first layer and/or the core.

The resultant core having the first layer thereon is prepared by any conventional means known in the pharmaceutical art, e.g. compression, tableting technology, spraying technology, or encapsulation in a pharmaceutically acceptable presentation, such as a gelatin capsule. In particular, typically the core formulation of the present invention is preferably fabricated by compression into a tablet.

The resultant core formulation of the present invention is useful to treat diabetes mellitus. Surprisingly the resultant core formulation of the invention is as user friendly and clinically effective as compared to the administration of metformin alone or troglitazone hydrochloride as demonstrated by co-administration of the two agents [*Whitcomb; et al., United States Patent No. 6,011,049*], where in general, the incidence of adverse events was not influenced by age or menopausal status, and further, patients treated with the combination therapy attained better glycemic control than with either monotherapy.

It is to be understood, however, that for any particular subject being treated, e.g., a mammal, specific dosage regimens should be adjusted according to the individual need. It is further to be understood that the dosages set forth herein are examples only and that they do not, to any extent limit the scope of the practice
5 of the present invention.

The core formulation of the present invention may be administered orally, for example, with inert diluent or with an edible carrier. For the purpose of oral therapeutic administration, the core formulation may have excipients incorporated therein. The subject core formulation may also contain the following
10 adjuvants: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel®, corn starch and the like; a lubricant such as magnesium stearate or Sterotex; a glidant such as colloidal silicon dioxide; and a sweetening agent such as sucrose or saccharin may be added or a flavoring agent such as peppermint, methyl
15 salicylate or orange flavoring.

The subject core formulation of the invention may contain other various materials which modify the physical form of the dosage unit (the subject core formulation), for example, as coatings. Thus, the subject core formulation of the present invention may be coated with sugar, shellac or other enteric coating
20 agents. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

In an alternative embodiment of the present invention. The resultant core formulation (having a first layer completely or partially covering the core), is treated whereby an outer shell, at least a portion of which comprises a biodegradable
25 material having a predetermined rate of degradation or metabolism in the host being treated, is formed which encloses the particles of the first layer and/or the core.

The biodegradable material is a high molecular weight compound, which is physiologically acceptable and decomposes in the body of the human being or other animal and is absorbed.

30 The biodegradable material, comprising the outer shell, having a predetermined rate of degradation or metabolism or break down, is selected from those materials well known in the art, which do not react with metformin and/or

troglitazone, e.g. its maleate, hydrochloride, etc. Such materials include, biodegradable polymers, such as polyorthoesters, polyanhydrides, polyamides based on glutamic acid, polyalkyl cyanoacrylates, polyesters of lactic and glycolic acid, polyactide polymers, cellulosic polymers, polyvinyl acetate, polyvinyl alcohol, polyvinylchloride, natural and synthetic rubbers, polyacrylates, polystyrene, etc. Additionally, reference is made to U.S. Patents Nos. 4,166,800, and 4,389,330, which disclose additional shell forming materials and are incorporated hereinto by reference in their entirety.

The shell encapsulating the particles of troglitazone, e.g. its hydrochloride, of the first layer and/or the particles of metformin of the core is obtained by any conventional microencapsulation process whereby microspheres of metformin and/or troglitazone, e.g. its hydrochloride, are formed, e.g. a solvent removal process, a phase separation technique, coacervation etc. In this regard reference is made to U.S. Patents Nos. 4,166,800 and 4,389,330, Conte et al, *J. Controlled Release*, vol. 26, (1993), pages 39-47; and U.S. Patent No. 4,839,177; all of which are incorporated hereinto by reference in their entirety.

In a variation of the above alternative embodiment, the resultant core formulation is treated whereby only the top surface area of the first layer comprising metformin has a shell coating thereon. In this regard, reference is made to U.S. Patent No. 5,916,584, incorporated hereinto by reference in its entirety, which describes the process for forming such a shell. The resulting core formulation having the first layer encapsulated by the shell comprising the shell material, is one which provides a delay time prior to release of the active ingredients, i.e. troglitazone and metformin, to the patient being treated for diabetes mellitus.

It is again to be understood that for either metformin or pioglitazone, any pharmaceutically acceptable form encompasses the free acids, free bases, salts and various hydrate forms, including semi-hydrate forms of these medicaments, as well as other pharmaceutical materials which are used in the formulation process as acceptable excipient materials generally known to those skilled in the art.

It is understood that any one of the biguanides, i.e. drugs having actions of stimulation of anaerobic glycolysis, is covered by this invention as these, like metformin, increase the sensitivity to insulin in peripheral tissues. These

compounds also are involved in the inhibition of glucose absorption from the intestine, suppression of hepatic gluconeogenesis, and inhibition of fatty acid oxidation. Examples of other typical biguanides included in this application are phenformin, buformin etc.

We claim:

1. A core formulation comprising,
 - (a) a first layer comprising troglitazone or a pharmaceutically acceptable salt thereof as an active ingredient,
 - 5 (b) a core, at least a portion of which is enclosed by said first layer, comprising a biguanide as an active ingredient.
2. The formulation as defined in claim 1 wherein said first layer comprises troglitazone hydrochloride present in an amount ranging from 1 mg to 15 mg and said core comprises metformin present in an amount ranging from 100 mg to
10 500 mg.
3. The formulation as defined in claim 1, which further comprises a biodegradable shell having a predetermined rate of degradation covering at least a portion of said first layer to provide a predetermined delay in the time period of release of at least said troglitazone active ingredient.
- 15 4. The formulation as defined in claim 1, wherein said troglitazone and/or said biguanide are present as biodegradable microspheres having a biodegradable shell coating and where said shell coating has a predetermined rate of degradation.
5. A method of administering troglitazone and metformin to a
20 mammal, which comprises treating the mammal with the formulation defined in claim 2.
6. A method for producing a controlled release formulation, which comprises:
 - (a) producing a hollow outer shell comprising a biodegradable
25 material having a predetermined rate of degradation to provide a predetermined delay in the time period of release of the contents destined to be enclosed by said shell;
 - (b) inserting a core comprising metformin and having an outer layer comprising troglitazone partially enclosing said core, into said hollow outer
30 shell; and
 - (c) sealing said core within said hollow outer shell.

7. A method of producing a combined formulation of troglitazone hydrochloride and metformin, which comprises:

- (a) forming a core of the metformin; and
- (b) depositing a layer of troglitazone hydrochloride on at least a
5 portion of a surface of said core.

8. A method of treating diabetes mellitus in a patient in need thereof, which comprises administering to the patient the formulation of claim 1 wherein said active ingredients are each present in an effective amount.

9. A pharmaceutical composition comprising an effective
10 amount of troglitazone hydrochloride combined with an effective amount of metformin.

10. A method of treating diabetes mellitus in a patient in need thereof, which comprises, administering to the patient the composition of claim 9.

11. A pharmaceutical composition comprising an effective
15 amount of troglitazone hydrochloride combined with an effective amount of phenformin.

12. A pharmaceutical composition comprising an effective amount of troglitazone hydrochloride combined with an effective amount of buformin.

20 13. A method of treating diabetes mellitus in a patient in need thereof, which comprises, administering to the patient the composition of claim 11.

14. A method of treating diabetes mellitus in a patient in need thereof, which comprises, administering to the patient the composition of claim 12.

25 15. A method of treating diabetes mellitus in a patient in need thereof, which comprises, administering to the patient the composition of claim 1 wherein the biguanide is phenformin.

16. A method of treating diabetes mellitus in a patient in need thereof, which comprises, administering to the patient the composition of claim 1 wherein the biguanide is buformin.

30 17. A method of treating diabetes mellitus in a patient in need thereof, which comprises, administering to the patient the composition of claim 1 wherein the biguanide is metformin.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/00039**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : A61K 9/32, 9/34, 9/36, 9/38

US CL : 424/474, 475, 477, 479, 480, 481, 482, 491, 493, 494, 496, 497

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/474, 475, 477, 479, 480, 481, 482, 491, 493, 494, 496, 497

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST, search terms: pioglitazone, troglitazone, biguanide, metformin, glucophage, rezulin, non insulin dependent diabetes, coated tablets, layered tablets, coated microcapsules

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category ^o	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6,011,049 A (WHITCOMB) 04 January 2000, see column 1, lines 41-50, column 4, line 42 through column 5, line 34, column 14, line 46 through column 15, claims 1-4 and 7-11.	1-17
Y	US 6,031,004 A (TIMMINS et al) 29 February 2000, see column 2, line 25 through column 3, column 5, lines 1-52, examples 5-8, claims 12-15.	1-17

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

07 FEBRUARY 2001

Date of mailing of the international search report

09 APR 2001

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JAMES M. SPEAR

Telephone No. (703) 308-1235

This Page Blank (uspto)